

SCHOTTKY BARRIER IN Au- AND Pd-CONTACTED SEMICONDUCTING CARBON NANOTUBES

*J. J. Palacios**, *P. Tarakeshwar***, and *Dae M. Kim***
jj.palacios@ua.es

*Departamento de Física Aplicada, Universidad de Alicante, San Vicente del Raspeig, Alicante 03690, Spain

**School of Computational Sciences, Korea Institute of Advanced Study, 207-43, Cheongyangni-2-dong, Dongdaemun-gu, Seoul 130-722, Korea

Work on semiconducting carbon nanotubes (CNT's) has intensified over the past few years in connection with their potential use as nanoscopic field-effect transistors [1]. Progress is, however, still hampered by the difficulties in understanding the contact between the metal electrodes and the CNT. Several groups are stepping up their efforts on the experimental side [2], progressively reaching a high degree of control on the contact formation and progressively gaining insight on this issue. To date, however, theoretical work lags behind, mostly focusing on developing simple microscopic models that can be handled at a reasonable computational cost [3], but with questionable prediction power.

Here we present work on the transport properties of semiconducting CNT coupled to commonly used metallic electrodes such as Au and Pd from a fully first-principles point of view. To this end we employ our code ALACANT (ALicante Ab initio Computation Applied to NanoTransport) [4]. The role of the chemistry at the contact, the geometry, and the atomic relaxation on the Schottky barrier and thus on the transport properties (n-type vs. p-type) is analyzed. Our preliminary results show that for simple but realistic contact geometries (see Figure 1) the Fermi level lies roughly in the middle of the gap for Au-contacted CNT's, while this one is pinned close to the top of the valence band for Pd-contacted CNT's. No appreciable band bending is observed for neither type of metal, although the charge transfer at the interface is larger in the case of Pd.




Figure 1. Relaxed structure of a semiconducting CNT contacted by the open ends to Pd electrodes.

References

- [1] J. Appenzeller, Y.-M. Lin, J. Knoch, and Ph. Avouris, *Phys. Rev. Lett.* **93**, 196805 (2004).
- [2] Harish M. Manohara, Eric W. Wong, Erich Schlecht, Brian D. Hunt, and Peter H. Siegel, *Nano Letters* (2006).
- [3] N. Nemeč, D. Tománek, and G. Cuniberti, *Phys. Rev. Lett.* **96**, 076802 (2006).
- [4] <http://www.guirisystems.com/alacant>.